

CANDIDATE NAME: _____

INDEX NUMBER _____



SERANGOON JUNIOR COLLEGE
JC2 PRELIMINARY EXAMINATION 2017

BIOLOGY 8875
Higher 1

CG _____

PAPER 1

Thursday
21 September 2017

1 hour

Additional materials:
Optical Mark Sheet

INSTRUCTIONS TO CANDIDATES

Write your name and CG in the spaces at the top of this page.

On the Optimal Mark Sheet, enter your name, subject title, test name, class. For your index number, enter your full NRIC number. Shade the corresponding lozenges on the OMS according to the instructions given by the invigilators.

AT THE END OF THE EXAMINATION, HAND IN BOTH THE OMS AND QUESTION PAPER.

INFORMATION FOR CANDIDATES

There are **thirty (30) questions** in this paper. Answer **all** questions. For each question, there are four possible answers, **A, B, C, D**. Choose the **one** you consider correct and record your choice in **soft pencil** on the OMS.

Each correct answer will score one mark. A mark will not be deducted for a wrong answer. Any rough working should be done on the question paper.


This question paper consists of 16 printed pages.

Answer all questions on the OTAS provided.

- 1 EDTA is used extensively as an anticoagulant for stored blood in blood banks. Thrombokinase plays a major role in the clotting of blood. EDTA decreases the reaction rate of thrombokinase by binding to calcium ions.

Which of the following describes the role of calcium ions?

- A Allosteric inhibitors
 - B Coenzymes
 - C Cofactors
 - D Competitive inhibitors
- 2 A cell in the G1 phase has two homologous pairs of chromosomes. It then undergoes two mitotic divisions. At the end of the second mitotic division, what is the total number of chromosomes and gene loci found in all the daughter cells formed?
- A 8 chromosomes and 4 times as many gene loci as the original parent cell.
 - B 8 chromosomes and 8 times as many gene loci as the original parent cell.
 - C 16 chromosomes and 4 times as many gene loci as the original parent cell.
 - D 16 chromosomes and 8 times as many gene loci as the original parent cell.
- 3 The electron micrograph below shows a liver cell.




Which statement(s) correctly describe(s) the labelled structures?

- 1 Structure **A** transports proteins from Structure **B** to Golgi Apparatus.
- 2 Proteins enter the lumen of Structure **B**, where they undergo chemical modifications such as glycosylation.
- 3 Structure **C** is starch grain.
- 4 The process shown in structure **D** is autolysis.

A 2 only **B** 1 and 2 only **C** 2 and 3 only **D** 2, 3 and 4 only


- 4 The graph represents the changes in the DNA content within a cell at different stages in the cell cycle.



Name the events occurring at **P**, **Q** and **R**, and identify the stage where meiosis is occurring.

	P	Q	R	Meiosis occurring at
A	S phase	Fertilisation	Cytokinesis	Y
B	Fertilisation	Interphase	Cytokinesis	Z
C	S phase	Prophase	Telophase	Y
D	Fertilisation	Metaphase	Telophase	Z

- 5 The graph shows changes in the amount of DNA in a cell during one cell cycle. The letters U – Z marks out the different phases in the cell cycle.



Many drugs that are used to treat cancer work at different time periods during the cell cycle.

- (i) Cisplatin binds to DNA and stops free DNA nucleotides from joining together.
- (ii) Drug B stops spindle fibres from shortening.


With reference to the cell cycle above, determine where these 2 drugs work.

	Cisplatin	Drug B
A	W	X
B	W	Y
C	U	X
D	U	Z

- 6 During the production of fruit juice, enzymes are used to break down the components of cell walls. Which carbohydrate will be produced by this hydrolysis?

- A Sucrose
- B Maltose
- C α - glucose
- D β - glucose

7 The figure below shows a DNA molecule.




Which statement(s) correctly describe the polynucleotide?

- 1 The structure labelled **A** corresponds to that of a purine, while the structure labelled **B** corresponds to that of a pyrimidine.
- 2 The antiparallel nature of DNA double helix allows phosphodiester bonds to form between the nitrogenous bases of opposite strands.
- 3 Distance between adjacent deoxyribonucleotides is 3.4 \AA and one turn consists of 10 deoxyribonucleotides. (Note: 10 \AA = 1 nm)
- 4 The wound DNA double helix consists of alternating major grooves and minor grooves along its axis which are essential for the binding with proteins.

- A** 1 only
B 1 and 2 only
C 3 and 4 only
D 1, 3 and 4 only

- 8 The RNA triplet UAG acts as a stop codon terminating the synthesis of a polypeptide. The diagram shows a template strand of DNA which codes for four amino acids.

Where would a mutation, introducing a thymine nucleotide, result in the termination of translation?



- 9 The non-template strand of a gene is analyzed and 20% of its bases are found to be adenine and 30% of its bases are cytosine. The corresponding template DNA strand of this gene has 10% cytosine.

What is the ratio of purine to pyrimidine found in pre-mRNA transcribed from this gene?

- A 1 : 1
- B 2 : 3
- C 3 : 2
- D 3 : 7
- 10 Which of the following statement(s) about cancer is / are true?
- I Individuals who inherit one mutant tumour suppressor gene are more likely to develop cancer than individuals with two non-mutant copies.
 - II Cancer is a result of increased cell division which promotes the mutation of a proto-oncogene.
 - III Mutagenic activation of a single oncogene is sufficient to cause a normal cell to develop into a cancerous cell.
- A I only
- B I and II only
- C I and III only
- D I, II and III

11 Which of the following statements correctly compares oxidative phosphorylation and non-cyclic photophosphorylation?

- A Both types of phosphorylation produce ATP and oxygen as end products.
- B Both types of phosphorylation produce ATP and the reduced form of a redox reagent.
- C Oxidative phosphorylation is involved in the conversion of one form of chemical energy to another while non-cyclic photophosphorylation is involved in converting light energy to chemical energy.
- D Water is an electron donor in non-cyclic photophosphorylation while it is an electron acceptor in oxidative phosphorylation.

12 What happens to most of the reduced NAD molecules in cell metabolism?

- A They act as oxidising agents in glycolysis.
- B They are oxidised in inner mitochondrial membrane for ATP formation.
- C They are oxidised in the Calvin cycle.
- D They combine with succinic acid as part of Krebs cycle.


13 Rotene and oligomycin are two metabolic poisons which affect cellular respiration. The effects of rotene and oligomycin on aerobic respiration are summarised in the table.

	Ability to use glucose	Ability to use oxygen	ATP yield
Rotene	Yes	No	Decreases
Oligomycin	Yes	Yes	Decreases

Which of the following correctly identifies the specific functions of these two metabolic poisons?

- | | Rotene | Oligomycin |
|----------|------------------------------|------------------------------|
| A | Electron transport inhibitor | Inhibits ATP synthase |
| B | Inhibits ATP synthase | Electron transport inhibitor |
| C | Dissipate proton gradient | Inhibits ATP synthase |
| D | Inhibits ATP synthase | Dissipate proton gradient |

- 14 In the graph below, the rate of CO₂ uptake by plant cells is shown to vary with increasing light intensity.




Which of the following is true at point X?


- A The plant is photosynthesizing.
 - B Rate of respiration equals rate of photosynthesis.
 - C CO₂ is a limiting factor.
 - D There is not enough light for photosynthesis to have commenced.
- 15 Which sequence of events correctly describes evolution?
- 1 Differential reproduction of the spiders occurs.
 - 2 A new selection pressure occurs.
 - 3 Allele frequencies within the spider population change.
 - 4 Poorly adapted spiders have decreased survivorship.

- A 2, 4, 1, 3 B 2, 4, 3, 1 C 4, 1, 3, 2 D 4, 3, 1, 2

- 16 The diagram shows a section through a chloroplast. Where would the products of photophosphorylation be used?



- 17 ^{14}C -labelled carbon dioxide was supplied to photosynthesising algae. The relative amounts of three organic compounds were measured. The diagram shows the results.



Which of the following are correct explanations for the graph above?

- I GP level falls as shown in graph 2 due to the absence of reduced NADP when light is are switched off.
- II GP level rises as shown in graph 1 due to the absence of ATP when light is switched off.
- III Levels of RuBP and GP are constant during periods of light as they serve as intermediates in the Calvin cycle.
- IV RuBP level falls as carboxylation of RuBP is independent of light as shown in graph 3.
- V Sucrose level falls as shown in graph 3 due to the absence of ATP and reduced NADP.

- A I, II and V only
- B I, II and III only
- C II, III and IV only
- D III, IV and V only


18 The following statements are some findings of scientists in an attempt to investigate the evolutionary relationship between the anteater, armadillo and pangolin.

- I Anteater, armadillo and pangolin feed primarily on insects such as ants.
- II Anteater, armadillo and pangolin have long tongue and strong digging limbs.
- III The tongues of the anteater and armadillo are connected to the hyoid bone while the tongue of pangolin is not.
- IV There is a higher percentage similarity between the DNA sequences of Anteater and armadillo than with the pangolin.
- V There is very low percentage similarity between the DNA sequences of anteater and pangolin as well as between the armadillo and pangolin.

Which of the following conclusions can be drawn from the statements given above?

- A The anteater and pangolin have experienced divergent evolution as shown by homologous structures between their hyoid bones and tongues.
- B The anteater and pangolin have experienced convergent evolution as shown by homologous structures in their hyoid bones and tongues.
- C The armadillo and pangolin have experienced divergent evolution as shown by the low similarity between their DNA sequences.
- D The anteater and armadillo have experienced divergent evolution as shown by similarities in their DNA sequences and homologous anatomical structures.

19 The pedigree chart below shows the inheritance of a recessive condition known as human albinism. Only homozygous recessive individuals are albinos.



What is the probability of individual 9 being a heterozygous carrier?

- A 0.00
- B 0.25
- C 0.50
- D 1.00


- 20 The feature of silky feathers in show fowl is caused by a recessive allele. A pure breeding bird with normal feathers was crossed with a bird with silky feathers and all the offspring were normal. The offspring were then allowed to interbreed.

Which of the following statements would be true about the F₂ generation?

- 1 The expected ratio of normal to silky would be 3:1.
- 2 Half of the F₂ birds would be heterozygous.
- 3 A quarter of the F₂ birds would be homozygous.
- 4 Some of the normal birds would be pure breeding.

- A 1, 2 and 3
B 2, 3 and 4
C 1, 2 and 4
D 1 and 2 only

- 21 During the process of polymerase chain reaction (PCR), the amount of DNA synthesised can be traced using fluorescent probes and the measurements are shown in the following plot. The process initially goes through an exponential phase followed by a plateau phase eventually.



Which of the following statements is **true**?

- A During the exponential phase, the number of DNA molecules synthesized after 15 cycles is 15².
- B During the exponential phase, the temperature is always maintained at the optimum temperature of 72°C hence there is rapid amplification.
- C During the plateau phase, the reaction mixture is being depleted of ribonucleotides.
- D During the plateau phase, *Taq* polymerase may be denatured.

- 22 The dashed lines in the template sequence represent a long sequence of bases to be amplified.


Template

5' ATTCGGACTTG ----- GTCCAGCTAGAGG 3'

3' TAAGCCTGAAC ----- CAGGTCGATCTCC 5'

Which of the following sets of primers can be used in the PCR for the amplification of the following DNA sequence?


- A 5' GTCCAGC 3' & 5' CCTGAAC 3'
 B 5' ATTCGGA 3' & 5' CCTCTAG 3'
 C 5' GGAAGTTG 3' & 5' GCTGGAC 3'
 D 5' AUUCGGA 3' & 5' GAUCUCC 3'
- 23 A family with a history of a genetic disease is studied using restriction digestion of the DNA samples containing the gene responsible for the disease. The pedigree chart of the family is aligned with the autoradiogram obtained from Southern blotting. (Shaded symbols in the pedigree chart indicate individuals affected by disease.)



Based on the information given, which of the following can be deduced?

- A The disease allele is dominant to the normal allele.
 B The mutation creates a new restriction site in the affected gene.
 C One of the parents in generation I is a carrier.
 D The offspring in generation II is a carrier.

- 24 Digestion of a 4 kb DNA molecule with *EcoRI* yields two fragments of 1 kb and 3 kb each. Digestion of the same molecule with *HindIII* yields fragments of 1.5 kb and 2.5 kb. Finally, digestion with *EcoRI* and *HindIII* in combination yields fragments of 0.5 kb, 1 kb and 2.5 kb. How would a restriction map indicating the positions of the *EcoRI* and *HindIII* cleavage sites look like?




- 25 Which of the following statements about the human genome project (HGP) is **false**?
- A HGP aims to identify all the genes in human and to determine the DNA sequences of these genes.
 - B HGP aims to allow genetic testing to take place for earlier detection of genetic diseases
 - C HGP allows defective genes to be replaced through gene therapy
 - D HGP allows comparative studies to be made between humans and other organisms to identify similar genes associated with diseases.
- 26 Recent advances in the field of stem cell research have shown that induced pluripotent stem cells (iPS cells) can be artificially derived from adult somatic cells. iPS cells are mostly similar to natural pluripotent cells. This implies that iPS cells can
- A theoretically differentiate into all cell types.
 - B theoretically differentiate into any of the three germ layers.
 - C theoretically differentiate into gametes.
 - D theoretically capable of transdifferentiation.

27 Which of the following regarding embryonic stem cells and hematopoietic stem cells is true?

- A As embryonic stem cells develop, they turned into hematopoietic stem cells as they lose their ability to differentiate into all types of cells.
- B Embryonic stem cells have more genes than hematopoietic stem cells and thus are able to form more types of cells.
- C Under normal conditions, embryonic stem cells express more of their genes compared to the hematopoietic stem cells.
- D Both stem cells are derived from the zygotic stem cells with the hematopoietic stem cells having a lower differentiation potential compared to the embryonic stem cells.

28 The pBR322 vector is used to clone a eukaryotic gene, which has been digested by the restriction endonuclease *BamHI*.




Following transformation, bacterial cells were grown in four different media, as shown below:

- 1 Nutrient broth containing ampicillin
- 2 Nutrient broth containing tetracycline
- 3 Nutrient broth containing ampicillin and tetracycline
- 4 Nutrient broth without ampicillin and tetracycline

Which of the following media would bacterial cells containing the recombinant plasmids grow in?

- A 4 only
- B 1 and 2
- C 2 and 3
- D 1 and 4

- 29 The diagram shows how genetically identical frogs can be developed from unfertilised frog eggs. The diploid number for frogs is 26.




Which combination of numbers correctly identifies the number of chromosomes in each type of cell?

	V	W	X
A	13	13	26
B	13	26	13
C	13	26	26
D	26	26	13

- 30** An attempt was made to produce Golden rice. To determine whether or not DNA from the daffodils and the bacterium had been successfully incorporated in the DNA of the rice, scientists used PCR and gel electrophoresis to produce DNA profiles.

The following DNA profiles belong to the original strain of rice, three strains **I** to **III** of genetically modified Golden rice, and the species of daffodil and bacterium used to incorporate beta-carotene genes in the rice.



Which one of the strain(s) of Golden rice has successfully incorporated DNA from both the daffodil and the bacterium?

- A** Strain **I** only
- B** Strain **II** only
- C** Strain **I** and **III** only
- D** Strain **II** and **III** only

End of Paper

CANDIDATE NAME: _____

INDEX NUMBER _____



SERANGOON JUNIOR COLLEGE
JC2 PRELIMINARY EXAMINATION 2017

BIOLOGY 8875
Higher 1

CG _____

PAPER 1

Thursday
21 September 2017

1 hour

Additional materials:
Optical Mark Sheet

INSTRUCTIONS TO CANDIDATES

Write your name and CG in the spaces at the top of this page.

On the Optimal Mark Sheet, enter your name, subject title, test name, class. For your index number, enter your full NRIC number. Shade the corresponding lozenges on the OMS according to the instructions given by the invigilators.

AT THE END OF THE EXAMINATION, HAND IN BOTH THE OMS AND QUESTION PAPER.

INFORMATION FOR CANDIDATES

There are **thirty (30) questions** in this paper. Answer **all** questions. For each question, there are four possible answers, **A, B, C, D**. Choose the **one** you consider correct and record your choice in **soft pencil** on the OMS.

Each correct answer will score one mark. A mark will not be deducted for a wrong answer. Any rough working should be done on the question paper.

This question paper consists of 16 printed pages.


Answer all questions on the OTAS provided.

- 1 EDTA is used extensively as an anticoagulant for stored blood in blood banks. Thrombokinase plays a major role in the clotting of blood. EDTA decreases the reaction rate of thrombokinase by binding to calcium ions.

Which of the following describes the role of calcium ions?

- A Allosteric inhibitors
 - B Coenzymes
 - C Cofactors**
 - D Competitive inhibitors
- 2 A cell in the G1 phase has two homologous pairs of chromosomes. It then undergoes two mitotic divisions. At the end of the second mitotic division, what is the total number of chromosomes and gene loci found in all the daughter cells formed?
- A 8 chromosomes and 4 times as many gene loci as the original parent cell.
 - B 8 chromosomes and 8 times as many gene loci as the original parent cell.
 - C 16 chromosomes and 4 times as many gene loci as the original parent cell.**
 - D 16 chromosomes and 8 times as many gene loci as the original parent cell.

- 3 The electron micrograph below shows a liver cell.




Which statement(s) correctly describe(s) the labelled structures?

- 1 Structure **A** transports proteins from Structure **B** to Golgi Apparatus.
- 2 Proteins enter the lumen of Structure **B**, where they undergo chemical modifications such as glycosylation.
- 3 Structure **C** is starch grain.
- 4 The process shown in structure **D** is autolysis.

A 2 only **B** 1 and 2 only **C** 2 and 3 only **D** 2, 3 and 4 only


- 4 The graph represents the changes in the DNA content within a cell at different stages in the cell cycle.



Name the events occurring at **P**, **Q** and **R**, and identify the stage where meiosis is occurring.

	P	Q	R	Meiosis occurring at
A	S phase	Fertilisation	Cytokinesis	Y
B	Fertilisation	Interphase	Cytokinesis	Z
C	S phase	Prophase	Telophase	Y
D	Fertilisation	Metaphase	Telophase	Z

- 5 The graph shows changes in the amount of DNA in a cell during one cell cycle. The letters U – Z marks out the different phases in the cell cycle.



Many drugs that are used to treat cancer work at different time periods during the cell cycle.

- (i) Cisplatin binds to DNA and stops free DNA nucleotides from joining together.
- (ii) Drug B stops spindle fibres from shortening.


With reference to the cell cycle above, determine where these 2 drugs work.

	Cisplatin	Drug B
A	W	X
B	W	Y
C	U	X
D	U	Z

- 6 During the production of fruit juice, enzymes are used to break down the components of cell walls. Which carbohydrate will be produced by this hydrolysis?

- A Sucrose
- B Maltose
- C α - glucose
- D β - glucose**

7 The figure below shows a DNA molecule.




Which statement(s) correctly describe the polynucleotide?

- 1 The structure labelled **A** corresponds to that of a purine, while the structure labelled **B** corresponds to that of a pyrimidine.
- 2 The antiparallel nature of DNA double helix allows phosphodiester bonds to form between the nitrogenous bases of opposite strands.
- 3 Distance between adjacent deoxyribonucleotides is 3.4 \AA and one turn consists of 10 deoxyribonucleotides. (Note: 10 \AA = 1 nm)
- 4 The wound DNA double helix consists of alternating major grooves and minor grooves along its axis which are essential for the binding with proteins.

- A** 1 only
B 1 and 2 only
C 3 and 4 only
D 1, 3 and 4 only

- 8 The RNA triplet UAG acts as a stop codon terminating the synthesis of a polypeptide. The diagram shows a template strand of DNA which codes for four amino acids.

Where would a mutation, introducing a thymine nucleotide, result in the termination of translation?



- 9 The non-template strand of a gene is analyzed and 20% of its bases are found to be adenine and 30% of its bases are cytosine. The corresponding template DNA strand of this gene has 10% cytosine.

What is the ratio of purine to pyrimidine found in pre-mRNA transcribed from this gene?

- A 1 : 1
- B 2 : 3
- C 3 : 2
- D 3 : 7
- 10 Which of the following statement(s) about cancer is / are true?
- I Individuals who inherit one mutant tumour suppressor gene are more likely to develop cancer than individuals with two non-mutant copies.
 - II Cancer is a result of increased cell division which promotes the mutation of a proto-oncogene.
 - III Mutagenic activation of a single oncogene is sufficient to cause a normal cell to develop into a cancerous cell.
- A I only
- B I and II only
- C I and III only
- D I, II and III

11 Which of the following statements correctly compares oxidative phosphorylation and non-cyclic photophosphorylation?

- A Both types of phosphorylation produce ATP and oxygen as end products.
- B Both types of phosphorylation produce ATP and the reduced form of a redox reagent.
- C Oxidative phosphorylation is involved in the conversion of one form of chemical energy to another while non-cyclic photophosphorylation is involved in converting light energy to chemical energy.**
- D Water is an electron donor in non-cyclic photophosphorylation while it is an electron acceptor in oxidative phosphorylation.

12 What happens to most of the reduced NAD molecules in cell metabolism?

- A They act as oxidising agents in glycolysis.
- B They are oxidised in inner mitochondrial membrane for ATP formation.**
- C They are oxidised in the Calvin cycle.
- D They combine with succinic acid as part of Krebs cycle.


13 Rotene and oligomycin are two metabolic poisons which affect cellular respiration. The effects of rotene and oligomycin on aerobic respiration are summarised in the table.

	Ability to use glucose	Ability to use oxygen	ATP yield
Rotene	Yes	No	Decreases
Oligomycin	Yes	Yes	Decreases

Which of the following correctly identifies the specific functions of these two metabolic poisons?

- | | Rotene | Oligomycin |
|----------|-------------------------------------|------------------------------|
| A | Electron transport inhibitor | Inhibits ATP synthase |
| B | Inhibits ATP synthase | Electron transport inhibitor |
| C | Dissipate proton gradient | Inhibits ATP synthase |
| D | Inhibits ATP synthase | Dissipate proton gradient |

- 14 In the graph below, the rate of CO₂ uptake by plant cells is shown to vary with increasing light intensity.




Which of the following is true at point X?


- A The plant is photosynthesizing.
 - B Rate of respiration equals rate of photosynthesis.
 - C CO₂ is a limiting factor.
 - D There is not enough light for photosynthesis to have commenced.
- 15 Which sequence of events correctly describes evolution?
- 1 Differential reproduction of the spiders occurs.
 - 2 A new selection pressure occurs.
 - 3 Allele frequencies within the spider population change.
 - 4 Poorly adapted spiders have decreased survivorship.

A 2, 4, 1, 3 B 2, 4, 3, 1 C 4, 1, 3, 2 D 4, 3, 1, 2

- 16 The diagram shows a section through a chloroplast. Where would the products of photophosphorylation be used? **A**



- 17 ^{14}C -labelled carbon dioxide was supplied to photosynthesising algae. The relative amounts of three organic compounds were measured. The diagram shows the results.



Which of the following are correct explanations for the graph above?

- I GP level falls as shown in graph 2 due to the absence of reduced NADP when light is are switched off.
- II GP level rises as shown in graph 1 due to the absence of ATP when light is switched off.
- III Levels of RuBP and GP are constant during periods of light as they serve as intermediates in the Calvin cycle.
- IV RuBP level falls as carboxylation of RuBP is independent of light as shown in graph 3.
- V Sucrose level falls as shown in graph 3 due to the absence of ATP and reduced NADP.

- A I, II and V only
- B I, II and III only
- C II, III and IV only**
- D III, IV and V only


18 The following statements are some findings of scientists in an attempt to investigate the evolutionary relationship between the anteater, armadillo and pangolin.

- I Anteater, armadillo and pangolin feed primarily on insects such as ants.
- II Anteater, armadillo and pangolin have long tongue and strong digging limbs.
- III The tongues of the anteater and armadillo are connected to the hyoid bone while the tongue of pangolin is not.
- IV There is a higher percentage similarity between the DNA sequences of Anteater and armadillo than with the pangolin.
- V There is very low percentage similarity between the DNA sequences of anteater and pangolin as well as between the armadillo and pangolin.

Which of the following conclusions can be drawn from the statements given above?

- A The anteater and pangolin have experienced divergent evolution as shown by homologous structures between their hyoid bones and tongues.
- B The anteater and pangolin have experienced convergent evolution as shown by homologous structures in their hyoid bones and tongues.
- C The armadillo and pangolin have experienced divergent evolution as shown by the low similarity between their DNA sequences.
- D The anteater and armadillo have experienced divergent evolution as shown by similarities in their DNA sequences and homologous anatomical structures.**

19 The pedigree chart below shows the inheritance of a recessive condition known as human albinism. Only homozygous recessive individuals are albinos.



What is the probability of individual 9 being a heterozygous carrier?


- A 0.00
- B 0.25
- C 0.50
- D 1.00**

- 20 The feature of silky feathers in show fowl is caused by a recessive allele. A pure breeding bird with normal feathers was crossed with a bird with silky feathers and all the offspring were normal. The offspring were then allowed to interbreed.

Which of the following statements would be true about the F₂ generation?

- 1 The expected ratio of normal to silky would be 3:1.
 - 2 Half of the F₂ birds would be heterozygous.
 - 3 A quarter of the F₂ birds would be homozygous.
 - 4 Some of the normal birds would be pure breeding.
- A 1, 2 and 3
- B 2, 3 and 4
- C 1, 2 and 4**
- D 1 and 2 only

- 21 During the process of polymerase chain reaction (PCR), the amount of DNA synthesised can be traced using fluorescent probes and the measurements are shown in the following plot. The process initially goes through an exponential phase followed by a plateau phase eventually.



Which of the following statements is **true**?

- A During the exponential phase, the number of DNA molecules synthesized after 15 cycles is 15².
- B During the exponential phase, the temperature is always maintained at the optimum temperature of 72°C hence there is rapid amplification.
- C During the plateau phase, the reaction mixture is being depleted of ribonucleotides.
- D During the plateau phase, Taq polymerase may be denatured.**

- 22 The dashed lines in the template sequence represent a long sequence of bases to be amplified.

Template


5' ATTCGGA CT TG ----- GTCCAGCTAGAGG 3'

3' TAAGCCTGAAC ----- CAGGTCGATCTCC 5'

Which of the following sets of primers can be used in the PCR for the amplification of the following DNA sequence?

- A 5' GTCCAGC 3' & 5' CCTGAAC 3'
- B 5' ATTCGGA 3' & 5' CCTCTAG 3'**
- C 5' GGA CT TG 3' & 5' GCTGGAC 3'
- D 5' AUUCGGA 3' & 5' GAUCUCC 3'


- 23 A family with a history of a genetic disease is studied using restriction digestion of the DNA samples containing the gene responsible for the disease. The pedigree chart of the family is aligned with the autoradiogram obtained from Southern blotting. (Shaded symbols in the pedigree chart indicate individuals affected by disease.)



Based on the information given, which of the following can be deduced?

- A The disease allele is dominant to the normal allele.**
- B The mutation creates a new restriction site in the affected gene.
- C One of the parents in generation I is a carrier.
- D The offspring in generation II is a carrier.

- 24 Digestion of a 4 kb DNA molecule with *EcoRI* yields two fragments of 1 kb and 3 kb each. Digestion of the same molecule with *HindIII* yields fragments of 1.5 kb and 2.5 kb. Finally, digestion with *EcoRI* and *HindIII* in combination yields fragments of 0.5 kb, 1 kb and 2.5 kb. How would a restriction map indicating the positions of the *EcoRI* and *HindIII* cleavage sites look like? **A**




- 25 Which of the following statements about the human genome project (HGP) is **false**?
- A HGP aims to identify all the genes in human and to determine the DNA sequences of these genes.
 - B HGP aims to allow genetic testing to take place for earlier detection of genetic diseases**
 - C HGP allows defective genes to be replaced through gene therapy
 - D HGP allows comparative studies to be made between humans and other organisms to identify similar genes associated with diseases.
- 26 Recent advances in the field of stem cell research have shown that induced pluripotent stem cells (iPS cells) can be artificially derived from adult somatic cells. iPS cells are mostly similar to natural pluripotent cells. This implies that iPS cells can
- A theoretically differentiate into all cell types.
 - B theoretically differentiate into any of the three germ layers.**
 - C theoretically differentiate into gametes.
 - D theoretically capable of transdifferentiation.

27 Which of the following regarding embryonic stem cells and hematopoietic stem cells is true?

- A As embryonic stem cells develop, they turned into hematopoietic stem cells as they lose their ability to differentiate into all types of cells.
- B Embryonic stem cells have more genes than hematopoietic stem cells and thus are able to form more types of cells.
- C Under normal conditions, embryonic stem cells express more of their genes compared to the hematopoietic stem cells.
- D Both stem cells are derived from the zygotic stem cells with the hematopoietic stem cells having a lower differentiation potential compared to the embryonic stem cells.**

28 The pBR322 vector is used to clone a eukaryotic gene, which has been digested by the restriction endonuclease *BamHI*.




Following transformation, bacterial cells were grown in four different media, as shown below:

- 1 Nutrient broth containing ampicillin
- 2 Nutrient broth containing tetracycline
- 3 Nutrient broth containing ampicillin and tetracycline
- 4 Nutrient broth without ampicillin and tetracycline

Which of the following media would bacterial cells containing the recombinant plasmids grow in?

- A 4 only
- B 1 and 2
- C 2 and 3
- D 1 and 4**

- 29 The diagram shows how genetically identical frogs can be developed from unfertilised frog eggs. The diploid number for frogs is 26.




Which combination of numbers correctly identifies the number of chromosomes in each type of cell?

	V	W	X
A	13	13	26
B	13	26	13
C	13	26	26
D	26	26	13

- 30 An attempt was made to produce Golden rice. To determine whether or not DNA from the daffodils and the bacterium had been successfully incorporated in the DNA of the rice, scientists used PCR and gel electrophoresis to produce DNA profiles.

The following DNA profiles belong to the original strain of rice, three strains I to III of genetically modified Golden rice, and the species of daffodil and bacterium used to incorporate beta-carotene genes in the rice.



Which one of the strain(s) of Golden rice has successfully incorporated DNA from both the daffodil and the bacterium?

- A Strain I only
- B Strain II only**
- C Strain I and III only
- D Strain II and III only

End of Paper

CANDIDATE NAME: _____

INDEX NUMBER _____

CG _____



SERANGOON JUNIOR COLLEGE
JC2 PRELIMINARY EXAMINATION 2017

BIOLOGY
Higher 1
8875

Tuesday
18 September 2017

2 hours

Additional materials:
Writing paper

READ THESE INSTRUCTIONS FIRST

Write your name and index number in the spaces at the top of this page and on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

For Section A answer **all** questions.

For Section B, answer only one question.

INFORMATION FOR CANDIDATES

The intended number of marks is given in brackets [] at the end of each question or part question.

FOR EXAMINER'S USE	
Paper 1 (MCQ)	/30
Paper 2	
Section A	
1	/10
2	/11
3	/10
4	/9
Total	/40
Section B	
6 or 7	/20
P2 Total	/60
TOTAL (P1+P2)	/90
TOTAL (100%)	/100

This question paper consists of 10 printed pages.

Section A [40 marks]

Answer **all questions** in the spaces provided.

Question 1

There have been many breakthroughs in stem cell research in recent years. It has been discovered that stem cells are involved in the replacement of worn-out cells and repair of damaged tissues. Further research is being conducted to better understand the mechanism involved in controlling the behaviour of stem cells in order to better manipulate them to treat various diseases and disorders.

- (a)** State the type of stem cells involved in the replacement of worn-out cells and repair of damaged tissues, and describe the unique properties of this type of stem cells. [2]

.....
.....
.....

Stem cells undergo cell division to produce genetically identical daughter cells. **Fig. 1.1** shows two cells, each at a different stage of cell division.




Fig. 1.1

- (b)(i)** With reference to **Fig. 1.1**, state the stages of cell division in **Cell A** and **Cell B**. [1]

Cell A:

Cell B:

- (ii) The dysregulation of cell cycle can result in cancer. Outline the checkpoints that are present in normal cells to prevent this from occurring. [2]

.....

.....

.....

Fig. 1.2 shows information about the movement of chromatids in a cell that has just started metaphase of mitosis.




Fig. 1.2

- (c)(i) With reference to Fig. 1.2, state the duration of metaphase in the cell. [1]

.....

- (ii) Complete line Y on the graph. [1]

(iii) Account for your answer in (c)(ii).

[3]

.....
.....
.....
.....

[Total: 10]

Question 2

A mutation was found in the gene coding for NADP oxidase in a family of flowering plant. NADP oxidase is an enzyme that converts NADPH to NADP⁺.

(a) Explain the role of NADPH in photosynthesis.

[2]

.....
.....
.....

(b) Using your knowledge of photosynthesis, predict the effect of this mutation on plants. [3]

.....
.....
.....
.....
.....
.....

Rubisco is an enzyme required in the light-independent stage of photosynthesis. **Fig. 2.1** shows the effect of increasing temperature on the activity of two variations of Rubisco, **Rubisco C** and **Rubisco S**.




Fig. 2.1

(c) With reference to **Fig. 2.1**, compare the effect of temperature on the two enzymes. [3]

.....

.....

.....

.....

(d) Explain how different alleles give rise to different Rubisco structure. [3]

.....

.....

.....

.....

[Total: 11]

Question 3

Fig. 3.1 shows the schematic representation of a series of protein complexes found on the inner membrane of organelle **X** present in brown adipocytes.




Fig. 3.1

(a)(i) State the identity of organelle **X**. [1]

.....

(ii) Outline how ATP is usually synthesised in the inner membrane of organelle **X**. [4]

.....
.....
.....
.....
.....
.....
.....

- (b) Brown adipocytes contain a unique protein, UCP1, which is not found in organelle **X** in any other cell type.

Evaluate the impact of UCP1 on ATP synthesis and suggest the physiological significance of brown adipose tissue. [3]

.....

.....

.....

.....

- (c) In other cell types, NADH and FADH₂ are used to drive ATP synthesis by ATP synthase. Using relevant information from **Fig. 3.1**, suggest and explain why more ATP is produced from NADH. [2]

.....

.....

.....

[Total: 10]

Question 4

The table below shows the amino acid differences in the cytochrome b protein between various vertebrates.

	Human	Elephant	Platypus	Ostrich	Starling	Crocodile	Lungfish	Coelacanth	Goldfish	Shark
Human		26	40	43	41	47	83	70	68	71
Elephant			45	45	48	50	84	72	63	74
Platypus				54	52	51	89	74	70	76
Ostrich					26	36	91	75	68	73
Starling						47	91	77	67	70
Crocodile							85	78	70	77
Lungfish								90	94	86
Coelacanth									83	78
Goldfish										88
Shark										

- (a) Explain how differences in amino acid sequences in the cytochrome b chain allow the establishment of the phylogenetic tree. [2]

.....

.....

.....

- (b) Suggest why homology still features prominently in evolutionary studies despite the advantages that molecular evidence can confer. [1]

.....

.....

Giant anteaters, armadillos and Australian numbats (*Myrmecobius fasciatus*) have many similar traits. This led some to believe that they were closely related.

Table 4.1 shows the comparison of four characteristics between the three mammals.

Mammal	Characteristics			
	Diet	Body	Snout	Tongue
Armadillo	Feed on insects	Covered by bony keratinised plates	Pointy snout	Long tongues
Giant Anteater	Feed on ants and termites	Covered by hair	Elongated narrow snout	Long tongues
Numbats	Feed on termites	Covered by hair	Narrow snout	Long tongues

(c) Explain why variation is important in selection. [2]

.....

.....

.....

(d) Explain how the evolution of long tongues in numbats supports Darwin's theory of natural selection. [4]

.....

.....

.....

.....

.....

.....

[Total: 9]

Section B [20 marks]

Answer **one** question in this section.

Write your answers on the separate writing paper provided.

Your answers may be illustrated by large, clearly labeled diagrams, ONLY where appropriate.

Your answers must be in continuous prose.

Question 5

- (a) Using the induced-fit hypothesis, explain the mode of action of enzymes. [6]
- (b) With reference to haemoglobin, explain the significance of bonds in maintaining the protein's structure and function. [8]
- (c) Discuss the social implications of genetically modifying plants. [6]

OR

Question 6

- (a) Compare competitive and non-competitive inhibition of enzyme action. [6]
- (b) Describe the process of mitosis and its importance in living cells. [8]
- (c) Discuss the ethical implications of genetically modifying plants. [6]

END OF PAPER

CANDIDATE NAME: _____

INDEX NUMBER _____

CG _____



SERANGOON JUNIOR COLLEGE
JC2 PRELIMINARY EXAMINATION 2017

BIOLOGY
Higher 1
8875

ANSWER SCHEME

2 hours

Additional materials:
Writing paper

READ THESE INSTRUCTIONS FIRST

Write your name and index number in the spaces at the top of this page and on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

For Section A answer **all** questions.

For Section B, answer only one question.

INFORMATION FOR CANDIDATES

The intended number of marks is given in brackets [] at the end of each question or part question.

FOR EXAMINER'S USE	
Paper 1 (MCQ)	/30
Paper 2	
Section A	
1	/10
2	/11
3	/10
4	/9
Total	/40
Section B	
6 or 7	/20
P2 Total	/60
TOTAL (P1+P2)	/90
TOTAL (100%)	/100

This question paper consists of 10 printed pages.

Section A [40 marks]

Answer **all questions** in the spaces provided.

Question 1

There have been many breakthroughs in stem cell research in recent years. It has been discovered that stem cells are involved in the replacement of worn-out cells and repair of damaged tissues. Further research is being conducted to better understand the mechanism involved in controlling the behaviour of stem cells in order to better manipulate them to treat various diseases and disorders.

(a) State the type of stem cells involved in the replacement of worn-out cells and repair of damaged tissues, and describe the unique properties of this type of stem cells. [2]

- **Adult stem cells [1]**

Any 2 properties [1]:

- **Undifferentiated cells found in differentiated tissues**
- **Multipotent → Able to differentiate into a limited range of cell types**
- **Able to undergo mitotic cell division for self-renewal**

Stem cells undergo cell division to produce genetically identical daughter cells. **Fig. 1.1** shows two cells, each at a different stage of cell division.




Fig. 1.1

(b)(i) With reference to **Fig. 1.1**, state the stages of cell division in **Cell A** and **Cell B**. [1]

Cell A: Prophase

Cell B: Anaphase

- (ii) The dysregulation of cell cycle can result in cancer. Outline the checkpoints that are present in normal cells to prevent this from occurring. [2]

Any 2

- **G₁ checkpoint:** assesses if the environmental conditions (presence of growth factors and nutrients, absence of DNA damage, adequate cell size) are favourable for cell division to proceed
- **G₂ checkpoint:** assesses if DNA replication is completed and cell size is adequate.
- **M checkpoint:** assesses if all chromosomes are attached to the mitotic spindle at their kinetochores and arrests the mitotic cell at metaphase if centromeres are not properly attached to kinetochore microtubules, hence preventing entry into anaphase.

Fig. 1.2 shows information about the movement of chromatids in a cell that has just started metaphase of mitosis.




Fig. 1.2

- (c)(i) With reference to Fig. 1.2, state the duration of metaphase in the cell. [1]

- **18 min**

(ii) Complete line Y on the graph. [1]

(iii) Account for your answer in (c)(ii). [3]

- Chromosomes align singly at the metaphase plate during metaphase of mitosis OR sister chromatids are attached to microtubules from opposite poles at metaphase
- Sister chromatids start to separate to become daughter chromosomes and migrate towards the opposite poles in anaphase, as shown at 18th min of line X when distance between chromatids starts to increase. Hence distance between chromatid and pole will start to decrease at 18th min.
- Distance between chromatids reach a plateau/maximum at 28th min, chromosomes arrived at opposite poles. Hence, distance between chromatid and pole will be minimum at 28th min.

[Total: 10]

Question 2

A mutation was found in the gene coding for NADP oxidase in a family of flowering plant. NADP oxidase is an enzyme that converts NADPH to NADP⁺.

(a) Explain the role of NADPH in photosynthesis. [2]

- Provides reducing power/H⁺ to reduce
- Phosphoglyceric acid (PGA)/glycerate-3-phosphate (GP) to glyceraldehyde-3-phosphate (GALP)/phosphoglyceraldehyde (PGAL)/triose phosphate (TP)

(b) Using your knowledge of photosynthesis, predict the effect of this mutation on plants. [3]

- Mutation will result in low or no NADP oxidase activity
- Less conversion of NADPH to NADP⁺ for light reaction of photosynthesis (OR less reduction of PGA to GALP in Calvin cycle)
- Less ATP is synthesized (or less glucose produced)

Rubisco is an enzyme required in the light-independent stage of photosynthesis. **Fig. 2.1** shows the effect of increasing temperature on the activity of two variations of Rubisco, **Rubisco C** and **Rubisco S**.




Fig. 2.1

(c) With reference to **Fig. 2.1**, compare the effect of temperature on the two enzymes. [3]

- Both Rubisco C and Rubisco S has an increased rate of reaction as temperature increases up to optimum temperature OR both Rubisco C and Rubisco S are denatured at temperatures higher than optimum.
- Rubisco C has a lower optimum temperature of 20°C as compared to Rubisco S at 50 °C where rate of reaction is at a maximum
- Rubisco C reaches a lower maximum rate of reaction of 5.5 a.u. at a faster rate as compared to Rubisco S which reaches a maximum rate of reaction of 6 a.u at a slower rate.

(d) Explain how different alleles give rise to different Rubisco structure. [3]

- Different alleles have different DNA nucleotide sequence that results in a different mRNA/codon sequence after transcription
- Thus will result in different amino acid sequence / primary structure after translation
- Different R group interactions between amino acids affects folding of the polypeptide chain, giving rise to different 3D conformation in the tertiary structure

[Total: 11]

Question 3

Fig. 3.1 shows the schematic representation of a series of protein complexes found on the inner membrane of organelle **X** present in brown adipocytes.




Fig. 3.1

(a)(i) State the identity of organelle **X**. [1]

Mitochondrion

(ii) Outline how ATP is usually synthesised in the inner membrane of organelle **X**. [4]

- **NADH and FADH₂ carry hydrogen in the form of protons and electrons where proton remain in the matrix and electrons are passed along the progressively lower energy electron carriers in the electron transport chain.**
- **Energy released is used to pump H⁺ from the matrix to the intermembrane space via active transport which sets up a concentration gradient (high H⁺ conc in intermembrane space, low conc in the matrix)**
- **H⁺ diffuse down the concentration gradient from the intermembrane space to matrix via the stalked particle**
- **Provides a proton motive force that drives the synthesis of ATP by ATP synthase by phosphorylation of ADP and inorganic phosphate (chemiosmosis).**

- (b) Brown adipocytes contain a unique protein, UCP1, which is not found in organelle **X** in any other cell type.

Evaluate the impact of UCP1 on ATP synthesis and suggest the physiological significance of brown adipose tissue. [3]

- **As UCP1 allows protons to leak back into the matrix without passing through the ATP synthase,**
- **Loss of H⁺ concentration gradient, no ATP will be synthesized**
- **The energy released from the spontaneous flow of protons through UCP1 is lost as heat, which helps to keep the organisms warm.**

- (c) In other cell types, NADH and FADH₂ are used to drive ATP synthesis by ATP synthase. Using relevant information from **Fig. 6.2**, suggest and explain why more ATP is produced from NADH. [2]

- **NADH donates electrons to complex I while FADH₂ donates to complex II. The energy released from transfer of electrons through the complexes is used to pump protons across the inner membrane.**
- **NADH allows for more chances to pumps more protons across the gradient, which powers the ATP synthase and gives us 3 ATP per molecule of NADH, while FADH₂ produces 2 ATP during the ETC because it gives up its electron to complex II, bypassing complex I.**

[Total: 10]

Question 4

The table below shows the amino acid differences in the cytochrome b protein between various vertebrates.

	Human	Elephant	Platypus	Ostrich	Starling	Crocodile	Lungfish	Coelacanth	Goldfish	Shark
Human		26	40	43	41	47	83	70	68	71
Elephant			45	45	48	50	84	72	63	74
Platypus				54	52	51	89	74	70	76
Ostrich					26	36	91	75	68	73
Starling						47	91	77	67	70
Crocodile							85	78	70	77
Lungfish								90	94	86
Coelacanth									83	78
Goldfish										88
Shark										

(a) Explain how differences in amino acid sequences in the cytochrome b chain allow the establishment of the phylogenetic tree. [2]

- Percentage of amino acid difference indicates relatedness where few difference indicates recent common ancestor
- Provides quantitative data to construct phylogenetic tree

(b) Suggest why homology still features prominently in evolutionary studies despite the advantages that molecular evidence can confer. [1]

Any 1

- Less expensive as it does not rely on machines
- DNA/protein samples might be limited or unavailable

Giant anteaters, armadillos and Australian numbats (*Myrmecobius fasciatus*) have many similar traits. This led some to believe that they were closely related.

Table 4.1 shows the comparison of four characteristics between the three mammals.

Mammal	Characteristics			
	Diet	Body	Snout	Tongue
Armadillo	Feed on insects	Covered by bony keratinised plates	Pointy snout	Long tongues
Giant Anteater	Feed on ants and termites	Covered by hair	Elongated narrow snout	Long tongues
Numbats	Feed on termites	Covered by hair	Narrow snout	Long tongues

(c) Explain why variation is important in selection. [2]

- Genetic variation results in variation in phenotype between individuals in a population
- Giving rise to different reproductive success between individuals OR
- Under a particular selective pressure, individuals with the advantageous variation are selected for.

(d) Explain how the evolution of long tongues in numbats supports Darwin's theory of natural selection. [4]

- Genetic variation give rise to different tongue lengths in (ancestral) numbats
- Under a selection pressure of limited food/ big termite nests/ deep termite nests/AVP
- Numbats with longer tongues have a selective advantage/ will be selected for because they can probe deeper into termite nests / AVP
- Numbats with longer tongues will survive, reproduce and pass down alleles encoding for longer tongues to the next generation
- Over time, the frequency of alleles coding for long tongues will increase.

[Total: 9]

Section B [20 marks]

Answer **one** question in this section.

Write your answers on the separate writing paper provided.

Your answers may be illustrated by large, clearly labeled diagrams, ONLY where appropriate.

Your answers must be in continuous prose.

Question 5

(a) Using the induced-fit hypothesis, explain the mode of action of enzymes. [6]

1. Enzyme specific in its action due to complementary 3D configuration/conformation of active site to that of substrate;
2. The induced fit model suggests that the enzyme and the substrate do not fit together exactly;
3. Effective collisions between enzymes and (specific) substrate molecules result in substrate binding to active site of enzyme;
4. The enzyme undergoes a 3D conformation change, which improves the fit between substrate and enzyme;
5. to form enzyme-substrate (ES) complexes;
6. Product formed that no longer fits into active site and is released;

(b) With reference to haemoglobin, explain the significance of bonds in maintaining the protein's structure and function. [8]

1. Peptide bonds between amine groups and carboxyl groups of amino acids at primary structure of organisation
2. Hydrogen bonds between –CO and –NH groups of the polypeptide backbone;
3. Ref. to overall 3D configuration/ globular shape of haemoglobin;
4. Each globin polypeptide is folded such that the bulk of the hydrophobic amino acid residues are buried in the interior of the globular structure;
5. Ref. to haem binding pocket lined with hydrophobic amino acids to provide a hydrophobic environment for hydrophobic haem group to bind;
6. Hydrophilic amino acid residues are on the outside;
7. Haemoglobin is soluble in aqueous medium and hence a good transport protein for oxygen in blood;
8. The two polypeptide chains in each dimer are held together by mainly hydrophobic interactions;
9. The two dimers are held together by weak hydrogen and ionic bonds;
10. Resulting in the ability of the two dimers to move with respect to each other;
11. This allows for cooperativity;
12. When an oxygen molecule binds to/is released from 1 haemoglobin subunit, the binding/ release induces a conformational change in the remaining subunit;
13. Which increases/ lowers the affinity for oxygen of the remaining three oxygen binding sites respectively;
14. This facilitates the loading and unloading of oxygen;

(c) Discuss the social implications of genetically modifying plants. [6]

- 1) Use of vectors which confer antibiotic-resistance might result in these genes being passed on to other potential harmful bacteria which hampers treatment.
- 2) New allergens produced that are dangerous to people with allergy who consume them.
- 3) Genetically modified crops might establish themselves as weeds as they are able to withstand unfavourable environmental conditions.
- 4) Spread of resistance from genetically modified crops to weeds might result in the production of superweeds that are resistant to herbicides.
- 5) Genetically engineered organisms, if introduced into the environment, might upset the balance of the ecosystem as it might lead to increased competition for space and nutrients.
- 6) Loss of biodiversity

OR

Question 6

(a) Compare competitive and non-competitive inhibition of enzyme action. [6]

(b)

Features	Competitive	Non-competitive
Structure of inhibitor	Resembles substrate;	Does not resemble substrate;
Binding site of inhibitor	Binds to active site of enzyme;	Binds to enzyme at a region other than the active site;
Mechanism of inhibition	Blocks substrates from binding to active site of the enzyme;	Blocks substrates from binding to active site by changing the conformation of the active site;
Effect of high substrate concentration on inhibition	Inhibition can be reversed at high substrate concentration;	Inhibition cannot be reversed at high substrate concentration;
Achieving V_{max}	V_{max} in the presence of inhibitor can be very close to that of reaction in the absence of inhibitor;	V_{max} in the presence of inhibitor is less than that of reaction in the absence of inhibitor;
Similarities:		
1. At low substrate concentration, rate of reaction in the presence of inhibitors is		

slower than that in the absence of inhibitor;

(c) Describe the process of mitosis and its importance in living cells.

[8]

Process of mitosis: any 6 from PMAT (at least one from each stage)

Prophase:

- **Chromatin condenses to chromosome**
- **Sister chromatids joined at centromere**
- **Centrioles migrate to opposite poles**
- **Mitotic spindles begin to form; nuclear envelope disintegrates**
- **Nucleolus disappears**

Metaphase:

- **Centrioles reached opposite poles**
- **Spindle fibres attached to kinetochore**
- **Chromosomes align on metaphase plate**

Anaphase

- **Centromeres divide**
- **Sister chromatids separate and move to opposite poles**
- **Shortening of kinetochore microtubules/spindle fibres**

Telophase

- **Chromosomes reached opposite poles**
- **Chromosomes uncoiled to chromatin fibers**
- **Spindle fibers disintegrates**
- **Nuclear envelope reforms**
- **Nucleoli reappears,**
- **Forms two genetically identical daughter nuclei**

Importance (max 2):

- Maintains **genetic stability** to produce **genetically identical** nuclei
- **Increase number of cells** for **growth**
- **Asexual reproduction**
- **Replace damaged cell** and **regeneration**

(d) Discuss the ethical implications of genetically modifying plants.

[6]

- **GM plants grown as crops may lead to consumers having allergies as foreign proteins are produced in the plants, companies need to label their GM crops for consumers to make informed choices / consumer safety is compromised;**
- **Animal genes may be introduced to plant genomes, leading to concern of vegetarians or some religious groups which followers are not allowed to consume certain animals;**
- **GM crops lead to benefits that rich countries can enjoy due to more financial resources at the expense of poorer countries (e.g. increasing dependence of poor countries on rich countries for expensive GM crops), increasing rich-poor divide;**

- GM crops can produce higher quality food to allow large companies that develop the technology / reduce costs to increase profit margins to out-compete small scale farmers, increasing rich-poor divide;
- Patenting the GM crops reduces them to the level of objects and if patenting is not allowed, a company might not be able to protect the results of their research program.
- Tampering with nature, where the mixing of genes among species may be seen as violation of organisms natural intrinsic values, crossing species boundaries;

END OF PAPER